

# Lidar Roughness Estimate for Vegetation within 2.5 m Above Terrain Metadata

*June 2024*



**ABMI** ALBERTA BIODIVERSITY  
MONITORING INSTITUTE



# Overview

## *Summary*

This dataset consists of files in GeoTiff format. The dataset is derived from lidar point cloud data. The dataset is intended to aid in the visualization and interpretation of point cloud data for use in landscape and vegetation analysis.

## *Description*

This dataset provides the roughness of the vegetation within 2.5 m above the ground. The raster provides information about the variation in height and roughness of the vegetation.

## *Methods*

The Roughness Estimate (R250) is produced using the LidR package<sup>1,2</sup> and dependencies in R. The Canopy Height Model is filtered using `nlas250` (las points below 2.5m) and height computed above ground using `pitfree(thresholds, max_edge)`. Grid resolution is set to 0.25. The roughness raster is calculated using `terrain()` function and `opt = c("roughness")`. The resulting R20 raster output is GeoTiff format with a file name derived from the original las file name appended with "\_R250.tif".

## *Credits*

This dataset includes products derived from lidar data collected and processed by the ABMI.

## *Acknowledgements*

We would like to acknowledge Brank Hricko, Stephanie Andrews, Amber Becker, John Simms and other ABMI staff, for the processing of lidar data to derivative files provided here. We would also like to acknowledge several funders who supported the project including the Government of Alberta and the Oil Sands Monitoring Program.

## *Contact Information*

If you have questions or concerns about the data, please contact:

Geospatial Centre  
Alberta Biodiversity Monitoring Institute  
CW 405 Biological Sciences Centre  
University of Alberta

---

<sup>1</sup> Roussel J, Auty D, Coops NC, Tompalski P, Goodbody TR, Meador AS, Bourdon J, de Boissieu F, Achim A (2020). "lidR: An R package for analysis of Airborne Laser Scanning (ALS) data." *Remote Sensing of Environment*, 251, 112061. ISSN 0034-4257, doi:10.1016/j.rse.2020.112061, <https://www.sciencedirect.com/science/article/pii/S0034425720304314>.

<sup>2</sup> Roussel J, Auty D (2023). *Airborne LiDAR Data Manipulation and Visualization for Forestry Applications*. R package version 4.0.3, <https://cran.r-project.org/package=lidR>.



Edmonton, Alberta, Canada, T6G 2E9  
Email: abmiinfo@ualberta.ca

### *Keywords*

LIDAR, Canopy Height Model, Roughness, Raster, Radius, Landscape Metric, LidR

### *Citation*

Alberta Biodiversity Monitoring Institute. Lidar Roughness Estimate for Vegetation within 2.5m Above Terrain Metadata (Version 1.4). Last modified June 10, 2024.

### *Use Limitations*

By accessing the Data, you agree to indemnify and hold harmless the ABMI and the ABMI's subsidiaries, affiliates, related parties, officers, directors, employees, agents, independent contractors, advertisers, partners, and co-branders, from any and all actions, proceedings, claims, demands, liabilities, losses, damages, and expenses which may be brought against or suffered by the ABMI or which it may sustain, pay or incur, arising or resulting from your violation of this clause.

The Data is provided on an "As Is" and "As Available" basis and the ABMI does not guarantee that the Data will be suitable for your purposes or requirements. Any interpretation of or reliance upon the Data is at your own cost and risk. The ABMI further states that the Data is subject to change, and the ABMI gives no guarantee that the content is complete, accurate, error or virus free, or up to date. The ABMI disclaims all warranties, conditions, and other terms of any kind, whether express or implied, whether in contract, tort (including liability for negligence) or otherwise, including, but not limited to any implied term of satisfactory quality, fitness for a particular purpose, non-infringement of intellectual property, and any standard of reasonable care and skill. While every effort has been made to ensure that the Data is complete, accurate, and current, the ABMI, their employees are not liable for any loss or damage arising directly or indirectly from the possession, publication, or use of, or reliance on, that information.

The ABMI recognizes the value of open data, data collaborations, and the efforts of our data collaborators and partners in the collection of the Data. For this reason, the Data is freely available for non-commercial use, including, but not limited to, research and educational purposes. No use of the Data may be made for resale without prior permission in writing from the ABMI.

## Data Product Specifications

### *Spatial Resolution*

The spatial resolution for R250: 0.25 metres

### *Processing Environment*

The processing environment to produce the GeoTiffs and shapefile is the R programming language, which includes R 4.2, Rtools 4.2 and RStudio Version:2023.06.0. The list of packages utilized includes LidR, raster, rgdal, sf, sp, spatial, and terra.

### *Resource Maintenance*

Resource maintenance update frequency: as needed



### *Spatial Reference*

Projected Coordinate System: NAD 1983 CSRS UTM Zone 11N  
Projection: Transverse Mercator  
WKID: 2955  
Authority: EPSG  
Linear unit: Metres (1.0)  
False Easting: 500000.0  
False Northing: 0.0  
Central Meridian: -117.0  
Scale Factor: 0.9996  
Latitude Of Origin: 0.0  
Geographic Coordinate System: NAD 1983 (CSRS)  
Angular Unit: Degree (0.0174532925199433)  
Datum: D North American 1983 CSRS  
Spheroid: GRS 1980  
Semimajor Axis: 6378137.0  
Semiminor Axis: 6356752.314140356  
Inverse Flattening: 298.257222101  
    NAD\_1983\_10TM\_AEP\_Forest  
    WKID: 3400 Authority: EPSG  
    Projection: Transverse Mercator  
    False Easting: 500000.0  
    False Northing: 0.0  
    Central Meridian: -115.0  
    Scale Factor: 0.9992  
    Latitude of Origin: 0.0  
    Linear Unit: Meter (1.0)  
    Geographic Coordinate System: GCS\_North\_American\_1983  
    Angular Unit: Degree (0.0174532925199433)  
    Prime Meridian: Greenwich (0.0)  
    Datum: D\_North\_American\_1983  
    Spheroid: GRS\_1980  
        Semi-major Axis: 6378137.0  
        Semi-minor Axis: 6356752.314140356  
        Inverse Flattening: 298.257222101

Projected Coordinate System: NAD 1983 CSRS UTM Zone 12N  
Projection: Transverse Mercator  
WKID: 2956  
Authority: EPSG  
Linear unit: Meters (1.0)  
False Easting: 500000.0  
False Northing: 0.0



Central Meridian: -111.0  
Scale Factor: 0.9996  
Latitude Of Origin: 0.0  
Geographic Coordinate System: NAD 1983 (CSRS)  
Angular Unit: Degree (0.0174532925199433)  
Datum: D North American 1983 CSRS  
Spheroid: GRS 1980  
Semimajor Axis: 6378137.0  
Semiminor Axis: 6356752.314140356  
Inverse Flattening: 298.257222101  
NAD\_1983\_10TM\_AEP\_Forest  
WKID: 3400 Authority: EPSG  
Projection: Transverse Mercator  
False Easting: 500000.0  
False Northing: 0.0  
Central Meridian: -115.0  
Scale Factor: 0.9992  
Latitude of Origin: 0.0  
Linear Unit: Meter (1.0)  
Geographic Coordinate System: GCS\_North\_American\_1983  
Angular Unit: Degree (0.0174532925199433)  
Prime Meridian: Greenwich (0.0)  
Datum: D\_North\_American\_1983  
Spheroid: GRS\_1980  
Semi-major Axis: 6378137.0  
Semi-minor Axis: 6356752.314140356  
Inverse Flattening: 298.257222101

### *Lineage*

The ABMI's Roughness Estimate for Vegetation within 2.5 m Above Terrain was built using ABMI collected lidar data. This dataset is divided into tiles and represents the roughness of the vegetation within 2.5 m above the ground on the landscape, as processed from the available lidar data.