

Projecting boreal bird responses to climate change: the signal exceeds the noise

D. STRALBERG,^{1,2,7} S. M. MATSUOKA,³ A. HAMANN,² E. M. BAYNE,¹ P. SÓLYMOS,¹ F. K. A. SCHMIEGELOW,^{2,4}
X. WANG,² S. G. CUMMING,⁵ AND S. J. SONG⁶

¹Department of Biological Sciences, CW 405, Biological Sciences Building, University of Alberta, Edmonton, Alberta T6G 2E9 Canada

²Department of Renewable Resources, 751 General Services Building, University of Alberta, Edmonton, Alberta T6G 2H1 Canada

³U.S. Fish and Wildlife Service, Migratory Bird Management, 1011 E. Tudor Road, Anchorage, Alaska 99503 USA

⁴Northern ENCS Program, Department of Renewable Resources, University of Alberta, c/o Yukon College, Whitehorse, Yukon Y1A 5K4 Canada

⁵Département des Sciences du Bois et de la Forêt, Pavillon Abitibi-Price, Bureau 2117, Université Laval, Québec, Québec G1V 0A6 Canada

⁶Canadian Wildlife Service, Environment Canada, 9250-49th Street, Edmonton, Alberta T6B 1K5 Canada

Abstract. For climate change projections to be useful, the magnitude of change must be understood relative to the magnitude of uncertainty in model predictions. We quantified the signal-to-noise ratio in projected distributional responses of boreal birds to climate change, and compared sources of uncertainty. Boosted regression tree models of abundance were generated for 80 boreal-breeding bird species using a comprehensive data set of standardized avian point counts (349 629 surveys at 122 202 unique locations) and 4-km climate, land use, and topographic data. For projected changes in abundance, we calculated signal-to-noise ratios and examined variance components related to choice of global climate model (GCM) and two sources of species distribution model (SDM) uncertainty: sampling error and variable selection. We also evaluated spatial, temporal, and interspecific variation in these sources of uncertainty. The mean signal-to-noise ratio across species increased over time to 2.87 by the end of the 21st century, with the signal greater than the noise for 88% of species. Across species, climate change represented the largest component (0.44) of variance in projected abundance change. Among sources of uncertainty evaluated, choice of GCM (mean variance component = 0.17) was most important for 66% of species, sampling error (mean = 0.12) for 29% of species, and variable selection (mean = 0.05) for 5% of species. Increasing the number of GCMs from four to 19 had minor effects on these results. The range of projected changes and uncertainty characteristics across species differed markedly, reinforcing the individuality of species' responses to climate change and the challenges of one-size-fits-all approaches to climate change adaptation. We discuss the usefulness of different conservation approaches depending on the strength of the climate change signal relative to the noise, as well as the dominant source of prediction uncertainty.

Key words: avian density; boosted regression trees; boreal birds; boreal forest; climate change; global climate models; signal-to-noise ratio; species distribution models; uncertainty; variance partitioning.

INTRODUCTION

Based on recent warming trends and global climate model (GCM) projections for the next century, the North American boreal forest is likely to experience particularly large changes in temperature and moisture availability (Balling et al. 1998, IPCC 2001). Climate change within the boreal region has already led to increased drought- and insect-induced tree mortality (Allen et al. 2010, Michaelian et al. 2010, Peng et al. 2011), wetland drying (Klein et al. 2005), and wildfire activity (Podur et al. 2002, Gillett et al. 2004, Soja et al. 2007). By virtue of its large size and relative intactness, the boreal forest is

thought to provide a large proportion of North America's breeding bird habitat (Wells and Blancher 2011). Therefore, species presently restricted to boreal regions may experience range reductions if those biomes shift northward and decrease in area, as projected for North America (Rehfeldt et al. 2012). However, positive temperature affinities and broad climatic tolerances suggest that many other species could expand their breeding distributions within the boreal region (Cumming et al. 2014). Recent northward range expansions of breeding birds have already been documented and attributed to climate change in temperate North America (Hitch and Leberg 2007), as well as in Europe (Thomas and Lennon 1999, Devictor et al. 2008).

As evidence has mounted for anthropogenic climate change and its widespread effects on species' distributions, it has increasingly been incorporated in systematic

Manuscript received 12 December 2013; revised 22 May 2014; accepted 17 June 2014. Corresponding Editor: J. Franklin.

⁷ E-mail: stralber@ualberta.ca

conservation planning efforts (e.g., Hannah et al. 2007, Carroll et al. 2009, Shaw et al. 2012). Accordingly, there is growing interest in forecasting the potential ecological impacts of climate change with an understanding of the associated uncertainties. Species distribution models (SDMs) have been widely used to project geographic changes in species' climatic habitat suitability (e.g., Peterson et al. 2002, Thuiller et al. 2005, Huntley et al. 2008). This correlative approach is based on the premise that the environmental niche of a species (*sensu* Grinnell 1917) may be quantified and used to develop spatial predictions of species' distributions under given environmental conditions, including future climates (Guisan and Zimmermann 2000, Peterson 2001). Paleo-ecological analysis of fossil pollen data suggests that, in the absence of climate conditions with no current analog, substitution of space for time is generally valid (Prentice et al. 1991, Huntley et al. 1993, Roberts and Hamann 2012), although differences in short-term predictability across taxa and ecological traits do exist (Kharouba et al. 2009, Dobrowski et al. 2011, Eskildsen et al. 2013). Passerine birds are not well represented in the fossil record, but molecular analyses are consistent with avian tracking of changes in climate and vegetation throughout Pleistocene glaciation cycles (Mengel 1964, Weir and Schluter 2004, Lovette 2005). Current climate has been found to be an important predictor of continental-scale avian distributions (Araújo et al. 2009, Jiménez-Valverde et al. 2011, Cumming et al. 2014), although not without skepticism (Bahn et al. 2006, Beale et al. 2008). Species' realized niches are also limited by biotic interactions (Hutchinson 1957), but empirical (Lovette and Hochachka 2006, Rubidge et al. 2011) and theoretical (Soberón 2007, Araújo and Rozenfeld 2014) evidence suggests that interactions that cannot be defined climatically are mostly local-scale processes that have minimal influence on broadscale distribution patterns (Rehfeldt et al. 2012).

Assuming that climatic niches of species are conserved over time (Wiens et al. 2010) and equilibrium with climate is maintained (Araújo and Pearson 2005), projecting species' long-term distributional responses to climate change will be problematic when uncertainty overpowers the prediction signal. Uncertainty surrounding future climate change trajectories (Murphy et al. 2004), combined with high variability among the SDMs themselves (Elith et al. 2006), has raised concerns about the utility and reliability of future projections. This has led to the development of ensemble forecasting approaches that use multiple models (Araújo and New 2007, Dormann et al. 2008*b*), as well as efforts to quantify and compare different aspects of prediction uncertainty. Although several studies have partitioned the variance in SDM-based future projections (Dormann et al. 2008*a*, Buisson et al. 2009, Diniz-Filho et al. 2009, Mbogga et al. 2010, Garcia et al. 2012), few have evaluated uncertainty with respect to the magnitude of predicted change (but see Thuiller 2004). High predic-

tion error may be outweighed by large directional changes in distribution and abundance. Thus, species-specific estimates of uncertainty ("noise") vs. change magnitude (the "signal") are needed over space and time.

A primary source of noise in future projections is the extrinsic variation among GCMs (hereafter "GCM uncertainty"). Although different GCMs are mostly based on the same physical principles (Jun et al. 2008, Masson and Knutti 2011, Pennell and Reichler 2011), the projections that they produce can be quite variable (Murphy et al. 2004, Kingston et al. 2009). Some GCMs are clearly better than others (Wang et al. 2007, Scherrer 2011), but metrics for model evaluation are not straightforward, and prediction patterns among GCMs can vary spatially as well as temporally (Tebaldi et al. 2005, Kang and Cressie 2013). Thus, the influence of GCM variability on SDM predictions depends not only on which model is considered, but also on the variables, seasons, and geographic areas that are important for a given species.

Considering the large variation among GCM projections, it is notable that intrinsic variation among SDM algorithms has often been found to be even larger (Dormann et al. 2008*a*, Buisson et al. 2009, Diniz-Filho et al. 2009, Garcia et al. 2012; but see Mbogga et al. 2010). However, high SDM variability may be driven by many factors, including the use of lower-performance algorithms (Elith et al. 2006, Hijmans and Graham 2006), sparse or inconsistent data (Araújo et al. 2005, Dormann et al. 2008*a*), poor or inconsistent model-building strategies (Meynard et al. 2013), extrapolation outside the range of data (Elith and Graham 2009), and improper handling of spatial dependence (Swanson et al. 2012). Some of these sources of error can be manipulated or controlled to reduce prediction uncertainty. In particular, by reducing model variability due to spatial dependence, inappropriate extrapolation, and model specification, one can focus on evaluating two fundamental sources of SDM-based (intrinsic) uncertainty: predictor variable selection and sampling error.

Predictor variable or model selection (hereafter "variable uncertainty") may have a large influence on SDM predictions (Mbogga et al. 2010, Synes and Osborne 2011, Braunisch et al. 2013), especially when important correlated variables decouple in the future. For example, many species' distributions are limited by the extent of agricultural land use (e.g., Siriwardena et al. 2000) or by the distribution of wetlands (e.g., Calmé and Desrochers 2000) more than by climate. Agriculture is constrained by current climate, particularly in northern environments, and the effects of climate and land use on bird distributions can be hard to disentangle (Clavero et al. 2011). However, land use will not necessarily track climate in the future, such that present confounding of climate and land use could lead to errors in future projections. Northern wetland distribution is also correlated with climate at continental extents, due




FIG. 1. Boreal and southern arctic study area shown in light gray, with boreal/arctic boundary as a gray line. Additional ecoregions projected to move into the study area by 2100 are depicted in darker gray. Point-count locations sampled for modeling are shown in black.

to the propensity for excess moisture to persist in colder environments. Despite reports of rapid boreal wetland drying attributed to recent climate change (e.g., Klein et al. 2005), loss of wetlands may take longer, depending on size and local hydrology, providing another mechanism by which presently correlated variables could become decoupled.

Sampling error (hereafter “sampling uncertainty”) results from various elements of epistemic uncertainty (“uncertainty associated with knowledge of the state of a system”; Regan et al. 2005), including measurement error, sampling bias, and inherent variability in the abundance of organisms across space and time (Elith et al. 2002). This type of uncertainty is often reflected in the differences among SDM algorithms that produce a wide range of individual model specifications. Multi-model predictions based on a diverse assortment of SDM techniques therefore can produce more robust predictions than single models (Thuiller et al. 2009). However, ensemble methods based on a single type of model (e.g., boosted regression trees) have similar strengths (Lawler et al. 2006), and bootstrapping methods may be used to estimate sampling error.

Here, we used an extensive boreal bird data set for North America (Cumming et al. 2010) to evaluate the

signal-to-noise ratio for projected changes in boreal bird abundance over the next century. We also compared different sources of uncertainty related to factors extrinsic (GCM uncertainty) and intrinsic (sampling and variable uncertainty) to SDMs. We evaluated spatial, temporal, and species-specific variation in each source of prediction uncertainty.

METHODS

Study area and avian survey data

We developed climate-change projections for boreal and southern arctic level II ecological regions as delineated by the Commission for Environmental Cooperation (CEC 1997). This included all subunits within the Taiga, Hudson Plain, and Northern Forests ecological regions, as well as the southern subunits of the Tundra ecological region (Alaska Tundra, Brooks Range Tundra, Southern Arctic), and boreal portion of the Northwestern Forested Mountains ecological region (Boreal Cordillera); see Fig. 1. We used data from avian point-count surveys (Ralph et al. 1995) that were conducted from 1992 to 2010 within the Nearctic boreal region (Brandt 2009), as compiled by the Boreal Avian Modelling (BAM) project (Cumming et al. 2010). This

included primarily off-road data from numerous inventory, monitoring, research, and impact assessment projects, including Provincial Breeding Bird Atlases and the Alaska Landbird Monitoring Survey, but also roadside point-count surveys conducted as part of the North American Breeding Bird Survey, BBS (Sauer et al. 2011). To account for anticipated shifts of southern climate conditions into currently boreal regions, we included point-count data from the same period collected in ecoregions within the continental United States and southern Canada that are south of the current boreal region, with climate conditions that are projected to shift northward into the study area within the next 100 years (Rehfeldt et al. 2012). This primarily consisted of BBS data, but also included off-road data from the western Great Lakes region (Hanowski and Niemi 1995).

Our initial compilation included data from 128 distinct projects with a total of 356 018 surveys at 125 547 unique locations (Fig. 1). To reduce the confounding influence of anthropogenic disturbance on modeled climate relationships, we removed surveys that had been conducted at agricultural, urban, or barren sites, according to the CEC's North American Land Change Monitoring System (NALCMS) land cover data set. We also removed surveys known to be conducted after recent timber harvest or other anthropogenic disturbance activities (not including fire), as mapped by Global Forest Watch Canada, the Alberta Biodiversity Monitoring Institute, and the United States LANDFIRE program. For our analysis, 349 629 surveys at 122 202 unique locations remained.

Climate data

Interpolated climate data were generated based on the parameter–elevation regressions on independent slopes model (PRISM) for the 1961–1990 normal period (Daly et al. 2008) and bioclimatic variables were derived according to Wang et al. (2012) and Hamann et al. (2013). Climate variables were chosen based on several criteria, including relevance to vegetation distributions (Hogg and Bernier 2005), avoidance of extreme colinearity (Dormann et al. 2013), and a preference for seasonal over annual variables when they showed high correlations. The final set of variables included extreme minimum temperature (EMT), chilling degree-days (DD0), growing degree-days (DD5), seasonal temperature difference (TD), mean summer precipitation (MSP), climate moisture index (CMI), and summer climate moisture index (CMIJJA). For complete variable definitions, see Appendix A: Table A1.

To represent potential future climates for three consecutive 30-year periods (2011–2040, 2041–2070, and 2071–2100), we used projections from the CMIP3 multi-model data set, corresponding to the fourth IPCC assessment report (Meehl et al. 2007). To limit computation time, we selected a subset of four complementary GCMs that spanned a range of project-

ed growing-season temperatures and precipitation levels within our study area: the German MPI ECHAM5, the Canadian CCCMA CGCM3.1, the United States GFDL CM2.1, and the United Kingdom Met Office HadGEM1 (Appendix A: Table A2, Fig. A1). Model projections were added as anomalies to the 4-km resolution 1961–1990 baseline data using the delta method and bilinear interpolation according to Wang et al. (2012). The data used in this study are part of a more comprehensive data set for North America, described in Appendix A and the Supplement. For this analysis, we adopted a scenario of high and monotonically increasing emissions (SRES A2; IPCC 2001), reflecting actual emissions during the decade elapsed since the scenario was defined (Raupach et al. 2007). Nineteen GCMs had runs available under the A2 emissions scenario.

The most highly correlated climate variables within our model-building data set (averaged across bootstrap samples, as described in the next section) were EMT and DD01 ($r = -0.88$), followed by MSP and CMIJJA ($r = -0.80$) (Appendix B: Table B1). Within the boreal and subarctic study area, the most highly correlated variables were CMI and CMIJJA ($r = 0.91$) and CMI and MSP ($r = 0.87$). Averaging across all 19 GCMs available for scenario A2, we found limited future decoupling (decrease in correlation over time) within the set of climate variables used, although there were differences between the model-building data set and prediction data sets (Appendix B: Fig. B1). EMT, DD0, and TD were the variables among which study area-wide correlations changed the most over time and were the most different from correlations within the model-building data set.

Land use and topography data

For a second set of models, we included a set of key land use/land cover variables that may influence bird abundance. We used the 250-m NALCMS land cover data set to calculate the current proportions of agriculture (AGRICULT), urban development (URBAN), open water (WATER), and wetlands (WETLAND) within each 4-km grid cell. We also derived a compound topographic index, CTI (Gessler et al. 1995)—or wetness index—from a 4-km digital elevation model and used it as a surrogate for wetland areas. The CTI was intended to differentiate lowland vs. upland vegetation types, in order to constrain future projections accordingly. These variables were not highly correlated with the seven climate indices or with each other (Appendix B: Table B1), and only minor future decoupling was observed.

We did not include proportions of natural upland land cover types, given the strong climatic basis for vegetation distribution at this resolution (Hamann and Wang 2006, McKenney et al. 2007). Limitation of data quality and coverage prevented adequate modeling and prediction of these remotely sensed land cover types relative to climate at such a broad scale.

Density models

We made use of the abundance information contained in point-count data by using survey- and species-specific correction factors, as described in Sólymos et al. (2013), to standardize density estimates across diverse protocols and environmental conditions. We examined 80 boreal-breeding passerine species with mapped breeding ranges (Ridgely et al. 2005) covering at least 10% of the boreal region (P. Blancher, *unpublished data*), as defined by Partners in Flight bird conservation regions 4, 6, 7, and 8 (Rich et al. 2004), which coincide with the boreal portion of our study area (Fig. 1). Each species also had surveys conducted with multiple time and/or distance intervals, general requirements for fitting the distance sampling (Buckland et al. 2001) and removal models (Sólymos et al. 2013) used to generate the correction factors.

We used boosted regression trees (BRT; De'ath 2007, Elith et al. 2008) to model avian densities at the level of the individual point-count station. We used the 'dismo' (Hijmans et al. 2011), 'gbm' (Ridgeway 2012) and 'raster' (Hijmans and van Etten 2012) packages for R (R Development Core Team 2012) to build BRT models for each species and then generate spatial predictions. We used the raw survey count at a point-count location as the response variable and included the log-transformed correction factors derived by Sólymos et al. (2013) as offsets to model avian density (number of males per hectare). For these count data, we specified a Poisson generalized boosted model (GBM) in the BRT estimation. The Poisson GBM uses an exponential function of the linear predictor within a gradient boosting algorithm (Friedman 2001, Ridgeway 2012).

We defined sampling units as the combination of the site (route, plot, or other local grouping of point counts) and 4-km grid cell ($n = 39\,186$ total sampling units); from each sampling unit with >10 surveys, we randomly selected a single point-count survey in each bootstrap iteration. This was to minimize spatial autocorrelation in surveys among points at the same site and temporal autocorrelation among surveys at the same point. We accounted for additional spatial autocorrelation among nearby sampling units by weighting the selection probabilities of each sampling unit by the inverse of the total number of surveys within the 20×20 km area surrounding the sampling unit. We minimized the influence of single data points by randomly selecting only one-third of the sampling units with ≤ 10 surveys in each bootstrap replicate. This procedure resulted in a total of 18 299 sampling units for each bootstrap replicate.

For each BRT model, we used a stepwise procedure and 10-fold cross-validation to identify the optimal number of trees needed to maximize the mean deviance explained. In each model run, we used a tree complexity of 3, learning rate of 0.001, and a bag fraction of 0.5. To ensure that the optimal number of trees could be found (Elith et al. 2008), we increased the learning rate to 0.005

if the limit of 10 000 trees was achieved, and reduced it to 0.0001 if fewer than 1000 trees were obtained. Using the optimal number of trees, we calculated 10-fold cross-validation statistics (proportion of deviance explained and Pearson's correlation coefficient) to assess prediction accuracy. For each species, we compared these statistics between the two variable sets (climate-only and climate + land use + topography) using paired t tests ($n = 11$ bootstrap replicates). Significance values were adjusted for multiple comparisons using a Holm (1979) correction. The importance of each model covariate was assessed by averaging the proportion of total deviance explained by a particular variable over all 11 bootstrap replicates.

Abundance projections

For each of the 80 boreal-breeding bird species, we applied the fitted BRT models to current and future climate conditions to predict avian density in each 4-km grid cell in the boreal/southern arctic study area. We multiplied the predicted density estimates for each grid cell (males/ha as an estimate of breeding pair density) by the grid cell area (1600 ha), and summed these values across grid cells to estimate total potential abundance.

We evaluated two sets of covariates: climate-only and climate + land use + topography. We generated models for 11 bootstrap samples, which were identical across species and covariate sets. We evaluated fitted BRTs under projected climates of alternate GCMs for three future time periods (2011–2040, 2041–2070, and 2071–2100).

For each species, we produced a total of 22 models from 11 bootstrap replicates and two variable sets for the 1961–1990 baseline period. We then produced a balanced set of future projections (264 total) for each species. This included predictions for all combinations of 11 bootstrap replicates, two variable sets, four GCMs, and three future time periods, for a total of 286 predictions for each species. To better assess the full range of variability across GCMs, we also generated an additional set of projections across the remaining 15 GCMs using just one bootstrap replicate, two variable sets, and three future time periods. This resulted in an additional 90 predictions, for a grand total of 376, including projections for the 1961–1990 baseline period.

Quantifying prediction uncertainty

To evaluate the overall signal-to-noise ratio for the projected change in overall abundance, we calculated Cohen's d (defined as the difference between means divided by the pooled standard deviation; Cohen 1992) for each species and each future time period compared to the baseline period, with variances pooled across the two time periods of interest for each calculation. For each time period, d was calculated for the full-factorial combination of 11 bootstrap replicates, two variable sets, and four GCMs ($n = 88$).

Next, to compare the climate change effect size to the variance components attributable to each source of uncertainty, we conducted two analyses of variance (ANOVA) for each species. Using projected change in overall abundance as the dependent variable, we first conducted four-factor ANOVAs with balanced data using just the four complementary GCMs ($n = 264$; 11 bootstrap replicates \times 2 variable sets \times 4 GCMs \times 3 future time periods). We partitioned the sums of squares among the effects of time period (climate change effect), sampling uncertainty, variable uncertainty, and GCM uncertainty, as well as the interactions between GCM and time, and variable and time. Variance components for each of these factors were calculated as the partial sum of squares divided by the total sum of squares (η^2). To evaluate the additional uncertainty introduced by considering the full suite of available GCMs (an additional 15 models), we also conducted an unbalanced ANOVA using type II sums of squares (Langsrud 2003) ($n = 354$; 264 original + 90 additional) with the 'car' (Fox and Weisberg 2011) package for R. For comparison across species, proportional abundance change was plotted against total uncertainty (sum of variance components for all uncertainty sources, including residuals).

Finally, to evaluate the relative magnitudes of each uncertainty source with respect to abundance projections, we calculated several versions of the coefficient of variation (CV); i.e., the standard deviation divided by the mean. For each species and time period, we calculated the CV in overall abundance for each source of uncertainty: sampling, variable, and GCM (future time periods only). Calculations were based on the full-factorial set of predictions (four complementary GCMs), and CV values for each uncertainty source were calculated with predictions for the other sources of uncertainty held constant at their average values. To evaluate spatial patterns of uncertainty for each species, using the same method, we also calculated the CV in density (males/ha) at the 4-km grid cell level for each source of uncertainty in each time period.

RESULTS

Model evaluation

All confidence ranges represent 5th and 95th percentiles except when otherwise noted. Across 80 species, prediction success of climate-only BRT models, assessed via cross-validation, averaged 0.222 (0.069, 0.462) in the deviance explained, and 0.225 (0.071, 0.474) in the Pearson correlation coefficient (Appendix C: Table C1). On average across species, the addition of land use and topographic variables to the climate-only models did not markedly improve cross-validation correlation (difference = 0.003 ± 0.014 SD) or deviance explained (difference = 0.002 ± 0.011 SD). However, for 18 of 80 species, the climate-only models were significantly improved by adding the land use and topography variables, in terms of one or both diagnostics after

multiple comparison correction. The climate-only model was significantly better for only one species.

Across species, temperature variables, on average, explained 0.145 (0.038, 0.330) of total deviance, and moisture variables explained 0.074 (0.017, 0.156) in climate-only models (Appendix C: Table C2). With models that also included land use and topographic variables, 0.051 (0.004, 0.167) of the deviance explained was accounted for by these additional variables, primarily agricultural land use proportion and compound topographic index (Appendix C: Table C3). Based on visual inspection of variable response curves from all 11 bootstrap runs, a total of 32 species exhibited clear monotonic decreases in abundance in response to agricultural land use proportion across bootstrap iterations; seven species had a clear negative response to urban land use proportion.

Projected changes in potential abundance

Of the 80 species modeled, 30 were projected to decline in potential abundance across the boreal and southern arctic regions by 2040; 34 species by 2070; and 37 species by 2100 (Appendix D: Table D1). Considering all sources of prediction uncertainty, projected declines were unequivocal (i.e., confidence intervals around projected change values did not contain zero) for 15 species by 2040, 18 by 2070, and 30 by 2100. Projected increases were unequivocal for 35 out of 50 species by 2040, 37 out of 46 by 2070, and 35 out of 43 by 2100. The distribution of projected species' responses shifted negatively and became increasingly dispersed over time, reflecting larger magnitudes of increase and decrease in abundance (Table 1).

Most species exhibited a northward (toward higher elevations) distributional shift in response to climate change (Appendix D: Fig. D1). Species' range centroids shifted an average of 18 m upward in elevation, 3 degrees north in latitude, and 3 degrees west in longitude by the end of the century (Table 2). Although areas of high richness of boreal species were projected to shift northward in distribution, total potential abundance across all species was projected to decline within the study area. Decreases in boreal species richness and density over time were most apparent in the interior west (Fig. 2). Projected current and future density layers (mean and CV across all sources of uncertainty) are *available online*.⁸

Prediction uncertainty relative to change

On average across 80 species, the signal-to-noise ratio, as measured by Cohen's d , was greater than 1 (i.e., signal > noise) for all three future time periods (Fig. 3). For the 2011–2040 time period, mean d was 1.42 (0.11, 3.49), and was greater than 1 for 46 species (Appendix D: Table D2). For the 2041–2070 time period, mean d

⁸ <http://borealbirds.databasin.org/>

TABLE 1. Frequency of projected percentage change in abundance across 80 boreal-breeding bird species.

Time period	Number of species showing the specified change in abundance					
	>50% decrease	25–50% decrease	25% decrease to 25% increase	25–50% increase	50–100% increase	>100% increase
2011–2040	0	7	50	18	4	1
2041–2070	2	10	33	7	16	10
2071–2100	10	16	17	6	10	21

Notes: Mean change values for each species are based on 11 bootstrap iterations, two variable sets, and four global climate models (GCMs). For individual species' projections, see Appendix D: Table D1.

increased to 2.38 (0.31, 5.32), and was greater than 1 for 59 species. By 2071–2100, mean d was 2.87 (0.73, 5.33), and was greater than 1 for 70 species.

Results from the balanced ANOVA indicated that the greatest source of variability in abundance predictions across species was attributed to the effect of climate change over time, with mean variance component = 0.442 (0.134, 0.760) (Table 3). When 19 GCMs were considered in an unbalanced ANOVA, the mean variance component of climate change decreased slightly to 0.397 (0.098, 0.725) (Table 3). For 21 species, sampling error represented the largest source of uncertainty (mean variance component = 0.118 across all 80 species) (Table 3; Appendix D: Table D3). For 13 of these species, sampling uncertainty was greater than the climate change effect. For 53 species, GCM represented the largest source of uncertainty (mean variance component = 0.174). However, it was only greater than the climate change effect for 10 species. When all 19 GCMs were considered, the mean variance component of GCM uncertainty increased to 0.228, compensating for a decrease in time- and sampling-related components. Variable selection resulted in a large variance component (up to 0.745) for a few species, but it was the greatest source of uncertainty for only four species (mean variance component = 0.047).

By definition, prediction uncertainty (calculated as the sum of all variance components except climate change from the balanced ANOVA) was negatively related to the magnitude of projected change in total abundance (Fig. 4), but a wide range of response magnitudes was seen along the range of prediction uncertainty. Prediction uncertainty was generally low relative to the projected magnitude of change for species with large projected increases. Species with the highest overall

prediction uncertainty were evenly split between those with high sampling uncertainty and those with high GCM uncertainty.

Spatial and temporal uncertainty

The magnitude and relative importance of the three components of prediction uncertainty changed over time (Table 4). Averaging across all 80 species, sampling error was the greatest source of uncertainty in current predictions of potential population size (as measured by CV), but uncertainty decreased over time, from an average of 0.129 (0.036, 0.390) in the current period to 0.099 (0.032, 0.228) by the end of the century. Variable-related uncertainty exhibited the opposite trend, increasing in importance over time from 0.058 (0.009, 0.163) to 0.115 (0.007, 0.347), as did uncertainty across the four GCMs, which more than doubled in magnitude from 0.092 (0.028, 0.161) in the 2011–2040 period to 0.216 (0.054, 0.505) in the 2071–2100 period (Table 4). When 19 GCMs were considered, the CV attributed to this component further increased to 0.266 (0.086, 0.630) by the end of the century.

Spatial patterns of uncertainty in the density predictions varied widely across species (Appendix D: Fig. D2), but for the current period it was concentrated in northern portions of the study area, where data are sparser (Fig. 5). Over time, areas of high sampling uncertainty were greatly reduced, as northern areas were projected to warm and thus more closely resemble the current climates of well-sampled boreal regions (Fig. 5a). By the end of the century, the small remaining areas of high variable uncertainty were concentrated in the western interior boreal region (Fig. 5b). Uncertainty based on four complementary GCMs increased over time, eventually overshadowing the other two sources of

TABLE 2. Projected changes over time in indices of spatial distribution (with 5th and 95th percentiles in parentheses) across 80 boreal-breeding bird species.

Time period	Mean latitude (°N)	Mean longitude (°W)	Mean elevation (m)
Current	56.2 (49.9, 63.3)	–98.7 (–115.7, –86.4)	473.2 (371.6, 686.8)
2011–2040	57.2 (50.3, 63.2)	–99.9 (–116.0, –88.0)	477.7 (375.1, 698.1)
2041–2070	58.0 (50.9, 64.1)	–100.8 (–116.3, –88.4)	481.4 (378.8, 674.5)
2071–2100	59.2 (52.4, 64.9)	–101.7 (–114.2, –89.3)	490.9 (387.5, 642.5)

Note: Species-level values are based on 11 bootstrap iterations, two variable sets, and four GCMs.




FIG. 2. Projected change over time in (a) boreal-breeding bird species richness and (b) total density of boreal-breeding birds. Species richness within each 4-km grid cell was calculated by converting density to probability of occurrence and summing probabilities across 80 species. The boreal/arctic boundary is shown in red.

uncertainty in most of the study area (Fig. 5c), with pockets of high GCM uncertainty concentrated in the northwest.

DISCUSSION

Signal vs. noise in projections of abundance

For projections of species' responses to future climate change to be useful, the magnitude of a species' projected response needs to be understood relative to the magnitude of uncertainty (Thuiller 2004). We found that for 58% of 80 boreal songbird species over the next 30 years—increasing to 88% of species by the end of the century—the climate change “signal” in

projections of abundance was greater than the “noise” generated by uncertainty due to a combination of sampling error, variable selection, and choice of global climate model (GCM). Despite the future increase in GCM uncertainty over time, this variability was swamped by the increasing magnitude of the projected directional change (positive or negative) in species abundance. This suggests that the predicted trajectories of avian responses to future climates are relatively robust for informing conservation planning and resource management decisions under climate change. Although the strength of a projected warming signal compared to GCM “noise” has been demonstrated (Kang and Cressie 2013), we found that the additional




FIG. 3. Change in the distribution of signal-to-noise ratios (Cohen's d) over time for 80 boreal bird species' abundance projections. The violin plots show the shapes of the distribution for each time period, with wider regions representing more common d values. Violin plots were generated using the beanplot package for R (Kampstra 2008) with a density bandwidth of 0.5 and Gaussian kernel estimation, where bandwidth is the standard deviation of the kernel. Gray lines link each species through the different time periods. The dashed line represents $d=1$; above this line, the climate change signal is greater than the noise due to the sources of uncertainty examined. Narrow black lines represent the 80 species; longer black lines extending beyond the violins represent mean d values for each time period.

uncertainties introduced by the species modeling process (specifically, sampling error and variable selection) generally did not overwhelm the climate change signal. This result appears consistent with Thuiller (2004), who found a majority (56%) consensus among different combinations of GCMs and SDMs for

mid-century species turnover projections in European plant communities.

Uncertainty due to GCM projections

By the end of the century, the largest source of prediction uncertainty across species was the choice of

TABLE 3. Variance components for four sources of variability (and interactions) in projected abundance change summarized across 80 boreal- and arctic-breeding species (a) for four complementary GCMs and (b) for all 19 GCMs available for the A2 emissions scenario (IPCC AR4).

Statistic	Time	Sampling	Variable	GCM	GCM \times time	Variable \times time	Remaining
a) Four GCMs							
Mean	0.442	0.118	0.047	0.174	0.107	0.017	0.095
2 SD	0.404	0.298	0.190	0.224	0.154	0.050	0.172
5th percentile	0.134	0.005	0.028	0.000	0.018	0.000	0.014
95th percentile	0.760	0.424	0.388	0.146	0.260	0.066	0.246
No. spp., largest	N/A	21	4	53	1	0	1
No. spp., >Time	N/A	13	3	10	6	0	9
b) All 19 GCMs							
Mean	0.397	0.090	0.045	0.228	0.134	0.016	0.090
2 SD	0.417	0.261	0.187	0.250	0.172	0.047	0.188
5th percentile	0.098	0.004	0.076	0.000	0.039	0.000	0.011
95th percentile	0.725	0.327	0.457	0.146	0.327	0.061	0.216
No. spp., largest	N/A	10	4	58	8	0	0
No. spp., >Time	N/A	13	3	21	14	0	11

Notes: Variance components are based on an ANOVA with three future time periods (climate change effect), two variable sets, and 11 bootstrap sampling iterations. No. spp., largest is the number of species for which that source of uncertainty was greatest; No. spp., >Time is the number of species for which that source of uncertainty was greater than the time effect. N/A means not applicable. The last column includes all other sources of variability, including unspecified interactions. For species-specific results, see Appendix D: Table D2.




FIG. 4. The magnitude of projected proportional change in abundance (y -axis) plotted against the magnitude of uncertainty (x -axis) by the end of the century (2071–2100) for 80 boreal bird species based on four complementary global climate models (GCMs). For species code identifications, see Appendix C: Table C1. Symbol fills indicate the greatest source of variability: climate change effect for species in gray; GCM for species in white; sampling or variable selection (CCSP only) for species in black. The x -axis represents the sum of all variance components except the climate change effect. The y -axis represents the log-transformed projected proportional change + 1 ($y = 0$ indicates no change).

GCM, indicating that multiple complementary GCMs should be used to capture the range of alternative futures. Direct comparison with other variance partitioning studies (Dormann et al. 2008a, Buisson et al. 2009, Diniz-Filho et al. 2009, Mbogga et al. 2010, Garcia et al. 2012) is complicated by differences in taxa, geographic regions, data resolution, and specific GCMs and time periods. However, our high-end estimate of 23% of variation due to GCM was comparable to the other study in this group (Garcia et al. 2012) that considered a full suite of available GCMs (17 vs. 19 in our study). The relatively small (6%) increase in variance from 4 to 19 GCMs reflects the high redundancy among these models (Masson and Knutti 2011) and suggests that a well-selected subset can appropriately reflect

climate model uncertainty. Furthermore, GCMs are not of equal accuracy (Scherrer 2011), so the use of poorly performing GCMs may be counterproductive (Räisänen 2007). Despite the large amount of uncertainty contributed by choice of GCM, we identified only 10 species for which the GCM-related uncertainty was consistently greater than the overall climate change effect (up to 21 when all 19 GCMs were considered). For these species primarily associated with deciduous forest (e.g., Mourning Warbler *Geothlypis philadelphia* and Canada Warbler *Cardellina canadensis*), future abundance trajectories were often nonlinear and diverged substantially over time, with larger projected decreases associated with the drier GCMs (Appendix D: Fig. D3). Areas of high GCM uncertainty were primarily located in the western interior

TABLE 4. Sources of prediction uncertainty (coefficient of variation) over time averaged across 80 boreal- and arctic-breeding bird species; confidence intervals in parentheses represent 5th and 95th percentiles.

Time period	Sampling	Variable	GCM-4	GCM-19
1961–1990	0.129 (0.036, 0.390)	0.058 (0.009, 0.163)	N/A	N/A
2011–2040	0.110 (0.028, 0.305)	0.051 (0.010, 0.136)	0.092 (0.028, 0.161)	0.131 (0.034, 0.276)
2041–2070	0.097 (0.027, 0.247)	0.072 (0.002, 0.230)	0.146 (0.041, 0.344)	0.184 (0.065, 0.385)
2071–2100	0.099 (0.032, 0.228)	0.115 (0.007, 0.347)	0.216 (0.054, 0.505)	0.266 (0.086, 0.630)

Notes: Sampling uncertainty is due to variation across 11 bootstrap samples; model uncertainty is due to variation between climate-only and climate + land use + topography models. Global climate model uncertainty is due to variation across four complementary models (GCM-4) and all 19 models (GCM-19) available for the A2 emissions scenario (IPCC AR4). N/A means not applicable.

boreal region, where available moisture is generally low, and small fluctuations in moisture may lead to major vegetation differences such as grassland vs. forest, or conifer vs. deciduous tree species (Hogg 1994, Schneider et al. 2009, Mbogga et al. 2010).

Uncertainty due to sampling error

Sampling error in the data used to build the model explained ~10% of the variance, on average, but much more for approximately a dozen species. The species with highest sampling error were generally less well represented in our data set, mostly due to their high-latitude affinities but also probably due to low densities; e.g., Rusty Blackbird *Euphagus carolinus* and American Pipit *Anthus rubescens*. This reflects the obvious fact that SDM accuracy may be reduced when limited occurrence data are available (Stockwell and Peterson 2002), especially when using more complex methods such as boosted regression trees (Wisiz et al. 2008). For this small subset of species with sparse data, current models probably could be improved by targeted surveys, especially in climatically underrepresented northern regions. However, despite large error bounds, most of these species were projected to decrease in abundance over time, with high signal-to-noise ratios. As such, their models are still informative, especially from the standpoint of identifying species most vulnerable to climate change. For almost all species examined, the sampling uncertainty decreased over time, as poorly sampled climates to the north were replaced by better-sampled climates to the south, i.e., eastern deciduous forest and interior grassland biomes. The relative lack of projected novel climate emergence (Williams et al. 2007) within our large study area (Rehfeldt et al. 2012) makes sampling uncertainty a much smaller problem than might otherwise be the case (e.g., Stralberg et al. 2009, Zurell et al. 2012). However, the regions with highest future sampling-related prediction uncertainty (mostly in Alaska) did tend to correspond with projected non-analog climates according to Rehfeldt et al. (2012), suggesting that signal-to-noise ratios may be much lower in regions that experience major novel climate development. These high-uncertainty regions may also be related to the partial decoupling of minimum annual temperature from temperature seasonality and growing-season heat sums (Appendix B).

Uncertainty due to predictor variables

Variable uncertainty was a minor component of the variability in future projections for all but a handful of the species that we evaluated. This probably reflected the low overall correlation between climate and land use variables in our data set, as well as the relatively strong predictive power of climate, compared to land use and topography, at a 4-km resolution. However, this source of uncertainty was important for a few species, primarily those with strong agricultural land use relationships such as Clay-colored Sparrow *Spizella pallida* (positive)

and Blue-headed *Vireo solitarius* (negative). This suggests that, when variable relationships are strong, minimal broadscale decoupling is sufficient for local variations in projections to arise. When climate and land use are confounded, the effects of climate on species' distributions may be overestimated, thereby misleading both the climate change projections (Clavero et al. 2011) and the conservation decisions based upon them. Consequently, there is a need for observational data sets that span a range of land use and climate conditions. This requirement is not always satisfied by roadside data from the North American Breeding Bird Survey (BBS; McKenney et al. 2001, Sauer et al. 2011), which comprise the primary distributional data available for climate change projection purposes in North America (e.g., Matthews et al. 2011; National Audubon Society 2014). In the boreal region in particular, roads and therefore BBS routes are simultaneously biased toward southern climates and agriculturally dominated landscapes (NABCI Canada 2012; Machtans et al. 2014). Our extensive data set, which included data from more remote parts of the boreal region, markedly reduced this bias.

Although the inclusion of land use and topography variables did not strongly influence range-wide predictions for most species, it was sometimes quite important locally. By the end of the century, variable uncertainty was concentrated in the southern portions of the boreal region, where the potential to support agricultural land uses in the future is greatest due to projected transition to prairie ecosystems (Frellich and Reich 2009). Unfortunately, boreal-wide spatially explicit projections of agricultural expansion generally do not exist, except for coarse (0.5° grid cell resolution) global projections that do not indicate any noticeable projected land-use conversion for the region (Strengers et al. 2004, Jetz et al. 2007, Hof et al. 2011). Other modeling efforts have focused on climatic suitability and plant hardiness zones (McKenney et al. 2001), which can be considered equivalent to our climate-only model and used to infer change (i.e., agricultural land uses shift with climate). However, projections that include socioeconomic and other policy drivers (e.g., Bierwagen et al. 2010, Radloff et al. 2011) would be necessary to adequately project local responses to the combined effects of climate and land-use change.

Other potential sources of uncertainty

It is rarely possible to evaluate and quantify all potential sources of uncertainty, some of which stem from vagueness of terms (Regan et al. 2005). For example, our models were intended to predict potential breeding-bird abundance based on climatic suitability. Actual bird numbers will depend on multiple demographic factors (e.g., overwinter survival and dispersal) that are not easily incorporated into a distribution modeling approach (but see Keith et al. 2008, Zurell et al. 2009, Fordham et al. 2013). Even within the more




FIG. 5. Spatial and temporal representations of the coefficient of variation in projected abundance at the 4-km grid cell level across multiple sources of uncertainty, averaged over 80 boreal-breeding bird species. (a) Sampling uncertainty due to variation across 11 bootstrap samples; (b) variable uncertainty due to variation between climate-only and climate + land use + topography models; and (c) GCM uncertainty due to variation across four complementary GCMs.

tangible sources of “epistemic” (known) uncertainty (Regan et al. 2005), we could not evaluate every potential contributor. Given the strong climatic basis for vegetation distribution at this resolution (Hamann and Wang 2006, McKenney et al. 2007), we assumed that avian responses to climate change would be driven by climate’s direct effects on vegetation, and we did not attempt to disentangle these effects. Thus the accuracy of short-term projections may be compromised when vegetation is in disequilibrium with climate (Svenning and Sandel 2013). However, because our intent was to evaluate trajectories of potential change rather than projected conditions for specific time periods, the

specific years were less important than the climate conditions that they represent.

Furthermore, our evaluation of variable uncertainty was limited to a subset of climate and land-use variables among which future decoupling was limited and localized. There may be other correlated, but unmeasured, climate variables, such as interannual variability (Cumming et al. 2014), that also decouple in the future, leading to additional prediction uncertainty. However, we were limited to existing GCM projections that do not yet adequately model changes to interannual climate variability (Mehta et al. 2010). More dramatically, inadequate representation of major positive feedbacks such as changes in albedo due to snow/ice (Screen and

Simmonds 2010) and cloud cover (Fasullo and Trenberth 2012) may also mean a substantial underestimation of climate sensitivity among current GCMs (Hansen et al. 2013). Consideration of more extreme scenarios could overwhelm the signal with noise, but we have focused here on generally accepted projections based on the so-called “fast feedbacks” (Rohling et al. 2012) for which short-term responses are better understood. Our results must be interpreted within these boundaries.

Conservation and management implications

A striking aspect of our results is the wide range of projected changes and uncertainty characteristics exhibited across species. Without quantitative analysis, we found some consistent and anticipated patterns among the species modeled. Species with the most northerly distributions often had high sampling uncertainty, due to sparse data, but also had large projected declines, leading to high signal-to-noise ratios that increased over time. Southern grassland-associated species were all projected to increase, but a combination of high variable uncertainty (due to positive associations with agriculture) and high GCM uncertainty led to low signal-to-noise ratios among this group. Deciduous forest-associated species tended to have high GCM uncertainty, given the potential for rapid broadscale conversion of deciduous and mixed forest to grassland, depending on “tipping points” in available moisture (Price et al. 2013). Although less common species generally had high sampling uncertainty, the signal-to-noise ratio for these species could easily be much higher than for common, high-abundance species with little projected response to climate change. Variability among species responses to climate change may be attributed to a variety of traits, the importance of which is not well understood (Kharouba et al. 2013). For birds, larger ranges are generally associated with lower model accuracy (Stockwell and Peterson 2002, McPherson et al. 2004, Segurado and Araújo 2004, McPherson and Jetz 2007). Other factors such as migratory behavior, trophic level/feeding guild, habitat specialization, and habitat association (especially wetland affinity) have been found to be important, but not consistently across regions and taxonomic subsets (Brotons et al. 2004, Huntley et al. 2004, Hernandez et al. 2006, McPherson and Jetz 2007). Further analysis is needed to better explain interspecific variation in climate change response, uncertainty, and signal-to-noise ratio.

However, this reinforces the individuality of species’ responses to climate change (Williams and Jackson 2007, Stralberg et al. 2009), and highlights the challenges of adopting one-size-fits-all approaches to climate change adaptation. Where feasible, land-based approaches that maintain natural disturbance dynamics (Noss 2001, Leroux et al. 2007) and facilitate broadscale distributional shifts, e.g., along gradients (Halpin 1997, Noss 2001, Hodgson et al. 2009), may prove most

effective in maintaining species diversity without requiring certainty about long-term outcomes. Such approaches are particularly viable in northern regions that are still relatively intact, such as the North American boreal forest.

However, individual species management is warranted for species of high conservation concern. Several studies have demonstrated the long-term inadequacy of relying solely on current environmental conditions to conserve and manage future species populations (Araújo et al. 2004, Veloz et al. 2013). Nevertheless, different conservation approaches may be justified depending on the strength of the climate change signal relative to the noise. When prediction uncertainty is high, there is greater risk associated with focusing on areas of predicted future climatic suitability (Fuller et al. 2008, Carroll et al. 2009, Carvalho et al. 2011). In these cases, a greater emphasis on areas of predicted overlap between current and future climatic distributions, i.e., macrorefugia (Keppel et al. 2012), may be appropriate. Conversely, justification is greater for an emphasis on future climate space when prediction uncertainty is low compared to the magnitude of change (Hamann and Aitken 2012, Oliver et al. 2012). The range of uncertainty exhibited across species also suggests a need for differential and quantitative weighting in assessments of climate change vulnerability (e.g., Gardali et al. 2012).

The dominant source of prediction uncertainty is also an important consideration in evaluating conservation and research strategies. For some species, especially those that are sensitive to changes in moisture balance, different GCMs result in distinctly different future trajectories. These species may be most effectively managed in an adaptive framework that considers the likelihood of alternative climate futures, updated as new information becomes available about GCM accuracy and reliability. Long-term monitoring at stationary locations will be a critical component of adaptive management efforts (Nielsen et al. 2009). For species with high sampling or variable uncertainty, the choice of GCM is less important within the already large range of future trajectories. In such cases, short-term efforts may be well spent by improving models through additional targeted sampling, e.g., in our case, in underrepresented northern regions, and in agricultural landscapes within marginal climates, respectively. It will also be important to study potential climate change effects on agricultural land uses (David and Marshall 2008), so that they may be factored into conservation decisions.

Finally, of immediate conservation concern within the boreal region is the rapid rate of industrial development, including forestry, energy, and other resource extraction, which could dramatically alter forest habitat over coming decades (Schneider et al. 2003, Hauer et al. 2010). Landscape-level effects of anthropogenic disturbance on avian communities (Schmiegelow et al. 1997, Drapeau et al. 2000, Hobson and Bayne 2000), and

avian vegetation type/age class relationships (Hobson and Schieck 1999, Schieck and Song 2006) have been identified regionally, and could be quantified across larger spatial extents with the development of comprehensive, standardized vegetation (e.g., Beaudoin et al. 2014; Cumming et al. *in press*) and anthropogenic disturbance (e.g., Pasher et al. 2013) data sets. With a better understanding of future development and vegetation trajectories, more temporally and spatially refined avian projections can also be generated. In the meantime, we suggest that bioclimatic models, when constructed carefully with accompanying uncertainty estimates, can provide useful projections for a majority of passerine species and should be interpreted in the context of associated uncertainties to inform conservation and management decisions.

ACKNOWLEDGMENTS

This publication is a contribution of the Boreal Avian Modelling (BAM) Project, an international research collaboration on the ecology, management, and conservation of boreal birds. We acknowledge the BAM Project's members, data partners, and funding agencies (including Environment Canada and the U.S. Fish and Wildlife Service). D. Stralberg was supported by a Vanier Canada Graduate Scholarship, as well as funding from the Climate Change and Emissions Management Corporation, the Alberta Ingenuity Fund, the Alberta Biodiversity Monitoring Institute, and the University of Alberta (Izaak Walton Killam Memorial Scholarship). We are grateful to Trish Fontaine for data management, to Rick Pelletier for computing resources, and to Nicole Barker and two anonymous reviewers for helpful feedback on earlier versions of this manuscript. Members of the Alberta Biodiversity Management and Climate Change Adaptation Project also provided insights valuable to this work.

LITERATURE CITED

- Allen, C. D., et al. 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259:660–684.
- Araújo, M. B., M. Cabeza, W. Thuiller, L. Hannah, and P. H. Williams. 2004. Would climate change drive species out of reserves? An assessment of existing reserve-selection methods. *Global Change Biology* 10:1618–1626.
- Araújo, M. B., and M. New. 2007. Ensemble forecasting of species distributions. *Trends in Ecology and Evolution* 22:42–47.
- Araújo, M. B., and R. G. Pearson. 2005. Equilibrium of species' distributions with climate. *Ecography* 28:693–695.
- Araújo, M. B., and A. Rozenfeld. 2014. The geographic scaling of biotic interactions. *Ecography* 37:406–415.
- Araújo, M. B., W. Thuiller, and N. G. Yoccoz. 2009. Reopening the climate envelope reveals macroscale associations with climate in European birds. *Proceedings of the National Academy of Sciences USA* 106:E45–E46.
- Araújo, M. B., R. J. Whittaker, R. J. Ladle, and M. Erhard. 2005. Reducing uncertainty in projections of extinction risk from climate change. *Global Ecology and Biogeography* 14:529–538.
- Bahn, V., R. J. O'Connor, and W. B. Krohn. 2006. Importance of spatial autocorrelation in modeling bird distributions at a continental scale. *Ecography* 29:835–844.
- Balling, R. C., P. J. Michaels, and P. C. Knappenberger. 1998. Analysis of winter and summer warming rates in gridded temperature time series. *Climate Research* 9:175–181.
- Beale, C. M., J. J. Lennon, and A. Gimona. 2008. Opening the climate envelope reveals no macroscale associations with climate in European birds. *Proceedings of the National Academy of Sciences USA* 105:14908–14912.
- Beaudoin, A., P. Y. Bernier, L. Guindon, P. Villemaire, X. J. Guo, G. Stinson, S. Magnussen, and R. J. Hall. 2014. Mapping attributes of Canada's forests at moderate resolution through kNN and MODIS imagery. *Canadian Journal of Forest Research* 44:521–532.
- Bierwagen, B. G., D. M. Theobald, C. R. Pyke, A. Choate, P. Groth, J. V. Thomas, and P. Morefield. 2010. National housing and impervious surface scenarios for integrated climate impact assessments. *Proceedings of the National Academy of Sciences USA* 107:20887–20892.
- Brandt, J. P. 2009. The extent of the North American boreal zone. *Environmental Reviews* 17:101–161.
- Braunisch, V., J. Coppes, R. Arlettaz, R. Suchant, H. Schmid, and K. Bollmann. 2013. Selecting from correlated climate variables: a major source of uncertainty for predicting species distributions under climate change. *Ecography* 36:971–983.
- Brotans, L., W. Thuiller, M. B. Araújo, and A. H. Hirzel. 2004. Presence-absence versus presence-only modelling methods for predicting bird habitat suitability. *Ecography* 27:437–448.
- Buckland, S. T., D. R. Anderson, K. P. Burnham, J. L. Laake, D. L. Borchers, and L. Thomas. 2001. *Introduction to distance sampling*. Oxford University Press, Oxford, UK.
- Buisson, L., W. Thuiller, N. Casajus, S. Lek, and G. Grenouillet. 2009. Uncertainty in ensemble forecasting of species distribution. *Global Change Biology* 16:1145–1157.
- Calmé, S., and A. Desrochers. 2000. Biogeographic aspects of the distribution of bird species breeding in Québec's peatlands. *Journal of Biogeography* 27:725–732.
- Carroll, C., J. R. Dunk, and A. Moilanen. 2009. Optimizing resiliency of reserve networks to climate change: multispecies conservation planning in the Pacific Northwest, USA. *Global Change Biology* 16:891–904.
- Carvalho, S. B., J. C. Brito, E. G. Crespo, M. E. Watts, and H. P. Possingham. 2011. Conservation planning under climate change: Toward accounting for uncertainty in predicted species distributions to increase confidence in conservation investments in space and time. *Biological Conservation* 144:2020–2030.
- CEC (Commission for Environmental Cooperation). 1997. *Ecological regions of North America: toward a common perspective*. Commission for Environmental Cooperation, Montreal, Canada.
- Clavero, M., D. Villero, and L. Brotans. 2011. Climate change or land use dynamics: do we know what climate change indicators indicate? *PLoS One* 6:e18581.
- Cohen, J. 1992. A power primer. *Psychological Bulletin* 112:155–159.
- Cumming, S. G., M. Houle, C. R. Drever, and P. Racine. *In press*. A gap analysis of tree species representation in protected areas within the Canadian boreal forest using digital forest resource inventory data. *Canadian Journal of Forest Research*.
- Cumming, S. G., K. L. Lefevre, E. Bayne, T. Fontaine, F. K. A. Schmiegelow, and S. J. Song. 2010. Toward conservation of Canada's boreal forest avifauna: design and application of ecological models at continental extents. *Avian Conservation and Ecology* 5:8.
- Cumming, S. G., D. Stralberg, K. L. Lefevre, P. Sólmos, E. M. Bayne, S. Fang, T. Fontaine, D. Mazerolle, F. K. A. Schmiegelow, and S. J. Song. 2014. Climate and vegetation hierarchically structure patterns of songbird distribution in the Canadian boreal region. *Ecography* 37:137–151.
- Daly, C., M. Halbleib, J. I. Smith, W. P. Gibson, M. K. Doggett, G. H. Taylor, J. Curtis, and P. P. Pasteris. 2008. Physiographically sensitive mapping of climatological temperature and precipitation across the conterminous United States. *International Journal of Climatology* 28:2031–2064.

- David, B. L., and B. B. Marshall. 2008. Why are agricultural impacts of climate change so uncertain? The importance of temperature relative to precipitation. *Environmental Research Letters* 3:034007.
- De'ath, G. 2007. Boosted trees for ecological modeling and prediction. *Ecology* 88:243–251.
- Devictor, V., R. Julliard, D. Couvet, and F. Jiguet. 2008. Birds are tracking climate warming, but not fast enough. *Proceedings of the Royal Society B* 275:2743.
- Diniz-Filho, J. A. F., L. M. Bini, T. F. Rangel, R. D. Loyola, C. Hof, D. Nogués-Bravo, and M. B. Araújo. 2009. Partitioning and mapping uncertainties in ensembles of forecasts of species turnover under climate change. *Ecography* 32:897–906.
- Dobrowski, S. Z., J. H. Thorne, J. A. Greenberg, H. D. Safford, A. R. Mynsberge, S. M. Crimmins, and A. K. Swanson. 2011. Modeling plant ranges over 75 years of climate change in California, USA: temporal transferability and species traits. *Ecological Monographs* 81:241–257.
- Dormann, C. F., et al. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46.
- Dormann, C. F., O. Purschke, J. R. G. Márquez, S. Lautenbach, and B. Schröder. 2008a. Components of uncertainty in species distribution analysis: a case study of the Great Gray Shrike. *Ecology* 89:3371–3386.
- Dormann, C. F., et al. 2008b. Prediction uncertainty of environmental change effects on temperate European biodiversity. *Ecology Letters* 11:235–244.
- Drapeau, P., A. Leduc, J.-F. Giroux, J.-P. L. Savard, Y. Bergeron, and W. L. Vickery. 2000. Landscape-scale disturbances and changes in bird communities of boreal mixed-wood forests. *Ecological Monographs* 70:423–444.
- Elith, J., M. A. Burgman, and H. M. Regan. 2002. Mapping epistemic uncertainties and vague concepts in predictions of species distribution. *Ecological Modelling* 157:313–329.
- Elith, J., and C. H. Graham. 2009. Do they? How do they? WHY do they differ? On finding reasons for differing performances of species distribution models. *Ecography* 32:66–77.
- Elith, J., et al. 2006. Novel methods improve prediction of species' distributions from occurrence data. *Ecography* 29:129–151.
- Elith, J., J. R. Leathwick, and T. Hastie. 2008. A working guide to boosted regression trees. *Journal of Animal Ecology* 77:802–813.
- Eskildsen, A., P. C. le Roux, R. K. Heikkinen, T. T. Høye, W. D. Kissling, J. Pöyry, M. S. Wisz, and M. Luoto. 2013. Testing species distribution models across space and time: high latitude butterflies and recent warming. *Global Ecology and Biogeography* 22:1293–1303.
- Fasullo, J. T., and K. E. Trenberth. 2012. A less cloudy future: the role of subtropical subsidence in climate sensitivity. *Science* 338:792–794.
- Fordham, D. A., et al. 2013. Population dynamics can be more important than physiological limits for determining range shifts under climate change. *Global Change Biology* 19:3224–3237.
- Fox, J., and S. Weisberg. 2011. *An R companion to applied regression*. Second edition. Sage, Thousand Oaks, California, USA.
- Frelich, L. E., and P. B. Reich. 2009. Will environmental changes reinforce the impact of global warming on the prairie-forest border of central North America? *Frontiers in Ecology and the Environment* 8:371–378.
- Friedman, J. H. 2001. Greedy function approximation: a gradient boosting machine. *Annals of Statistics* 29:1189–1232.
- Fuller, T., D. P. Morton, and S. Sarkar. 2008. Incorporating uncertainty about species' potential distributions under climate change into the selection of conservation areas with a case study from the Arctic Coastal Plain of Alaska. *Biological Conservation* 141:1547–1559.
- Garcia, R. A., N. D. Burgess, M. Cabeza, C. Rahbek, and M. B. Araújo. 2012. Exploring consensus in 21st century projections of climatically suitable areas for African vertebrates. *Global Change Biology* 18:1253–1269.
- Gardali, T., N. E. Seavy, R. T. DiGaudio, and L. A. Comrack. 2012. A climate change vulnerability assessment of California's at-risk birds. *PLoS One* 7:e29507.
- Gessler, P. E., I. D. Moore, N. J. McKenzie, and P. J. Ryan. 1995. Soil-landscape modelling and spatial prediction of soil attributes. *International Journal of Geographical Information* 9:421–432.
- Gillett, N. P., A. J. Weaver, F. W. Zwiers, and M. D. Flannigan. 2004. Detecting the effect of climate change on Canadian forest fires. *Geophysical Research Letters* 31:L18211.
- Grinnell, J. 1917. The niche-relationships of the California Thrasher. *Auk* 34:427–434.
- Guisan, A., and N. E. Zimmermann. 2000. Predictive habitat distribution models in ecology. *Ecological Modelling* 135:147–186.
- Halpin, P. N. 1997. Global climate change and natural-area protection: management responses and research directions. *Ecological Applications* 7:828–843.
- Hamann, A., and S. N. Aitken. 2012. Conservation planning under climate change: accounting for adaptive potential and migration capacity in species distribution models. *Diversity and Distributions* 19:268–280.
- Hamann, A., and T. Wang. 2006. Potential effects of climate change on ecosystem and tree species distribution in British Columbia. *Ecology* 87:2773–2786.
- Hamann, A., T. Wang, D. L. Spittlehouse, and T. Q. Murdock. 2013. A comprehensive, high-resolution database of historical and projected climate surfaces for western North America. *Bulletin of the American Meteorological Society* 94:1307–1309.
- Hannah, L., G. Midgley, S. Anelman, M. Araújo, G. Hughes, E. Martinez-Meyer, R. Pearson, and P. Williams. 2007. Protected area needs in a changing climate. *Frontiers in Ecology and the Environment* 5:131–138.
- Hanowski, J. M., and G. J. Niemi. 1995. Experimental design considerations for establishing an off-road, habitat-specific bird monitoring program using point-counts. Pages 145–150 in C. J. Ralph, J. R. Sauer, and S. Droege, editors. *Monitoring bird populations by point counts*. USDA Forest Service General Technical Report PSW-GTR-149. USDA Forest Service, Pacific Southwest Research Station, Albany, California, USA.
- Hansen, J., P., et al. 2013. Assessing “dangerous climate change”: required reduction of carbon emissions to protect young people, future generations and nature. *PLoS One* 8:e81648.
- Hauer, G., S. Cumming, F. Schmiegelow, W. Adamowicz, M. Weber, and R. Jagodzinski. 2010. Tradeoffs between forestry resource and conservation values under alternate policy regimes: A spatial analysis of the western Canadian boreal plains. *Ecological Modelling* 221:2590–2603.
- Hernandez, P. A., C. H. Graham, L. L. Master, and D. L. Albert. 2006. The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography* 29:773–785.
- Hijmans, R. J., and C. H. Graham. 2006. The ability of climate envelope models to predict the effect of climate change on species distributions. *Global Change Biology* 12:2272–2281.
- Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith. 2011. *Package dismo: species distribution modeling*. Version 0.7-23. R Foundation for Statistical Computing, Vienna, Austria. <http://cran.r-project.org/web/packages/dismo/index.html>
- Hijmans, R. J., and J. van Etten. 2012. *Package raster: geographic data analysis and modeling*. Version: 1.9-55. R

- Foundation for Statistical Computing, Vienna, Austria. <http://cran.r-project.org/web/packages/raster/index.html>
- Hitch, A. T., and P. L. Leberg. 2007. Breeding distributions of North American bird species moving north as a result of climate change. *Conservation Biology* 21:534–539.
- Hobson, K. A., and E. Bayne. 2000. Breeding bird communities in boreal forests of western Canada: consequences of “unmixing” the mixedwoods. *Condor* 102:759–769.
- Hobson, K. A., and J. Schieck. 1999. Changes in bird communities in boreal mixedwood forest: harvest and wildfire effects over 30 years. *Ecological Applications* 9:849–863.
- Hodgson, J. A., C. D. Thomas, B. A. Wintle, and A. Moilanen. 2009. Climate change, connectivity and conservation decision making: back to basics. *Journal of Applied Ecology* 46:964–969.
- Hof, C., M. B. Araújo, W. Jetz, and C. Rahbek. 2011. Additive threats from pathogens, climate and land-use change for global amphibian diversity. *Nature* 480:516–519.
- Hogg, E. H. 1994. Climate and the southern limit of the western Canadian boreal forest. *Canadian Journal of Forest Research* 24:1835–1845.
- Hogg, E. H., and P. Y. Bernier. 2005. Climate change impacts on drought-prone forests in western Canada. *Forestry Chronicle* 81:675–682.
- Holm, S. 1979. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics* 6:65–70.
- Huntley, B., Y. C. Collingham, S. G. Willis, and R. E. Green. 2008. Potential impacts of climatic change on European breeding birds. *PLoS One* 3:e1439.
- Huntley, B., R. E. Green, Y. C. Collingham, J. K. Hill, S. G. Willis, P. J. Bartlein, W. Cramer, W. J. M. Hageheijer, and C. J. Thomas. 2004. The performance of models relating species geographical distributions to climate is independent of trophic level. *Ecology Letters* 7:417–426.
- Huntley, B., R. A. Spicer, W. G. Chaloner, and E. A. Jarzembowski. 1993. The use of climate response surfaces to reconstruct palaeoclimate from Quaternary pollen and plant macrofossil data [and discussion]. *Philosophical Transactions of the Royal Society B* 341:215–224.
- Hutchinson, G. E. 1957. Concluding remarks. *Cold Spring Harbor Symposium on Quantitative Biology* 22:415–427.
- IPCC. 2001. Climate change 2001: the scientific basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK.
- Jetz, W., D. Wilcove, and A. Dobson. 2007. Projected impacts of climate and land-use change on the global diversity of birds. *PLoS Biology* 5:e157.
- Jiménez-Valverde, A., N. Barve, A. Lira-Noriega, S. P. Maher, Y. Nakazawa, M. Papeş, J. Soberón, J. Sukumaran, and A. Townsend Peterson. 2011. Dominant climate influences on North American bird distributions. *Global Ecology and Biogeography* 20:114–118.
- Jun, M., R. Knutti, and D. W. Nychka. 2008. Spatial analysis to quantify numerical model bias and dependence. *Journal of the American Statistical Association* 103:934–947.
- Kampstra, P. 2008. Beanplot: a boxplot alternative for visual comparison of distributions. *Journal of Statistical Software* 28:1–9. <http://www.jstatsoft.org/v28/c01/>
- Kang, E. L., and N. Cressie. 2013. Bayesian hierarchical ANOVA of regional climate-change projections from NARCCAP phase II. *International Journal of Applied Earth Observation and Geoinformation* 22:3–15.
- Keith, D. A., H. R. Akçakaya, W. Thuiller, G. F. Midgley, R. G. Pearson, S. J. Phillips, H. M. Regan, M. B. Araújo, and T. G. Rebelo. 2008. Predicting extinction risks under climate change: coupling stochastic population models with dynamic bioclimatic habitat models. *Biology Letters* 4:560–563.
- Keppel, G., K. P. Van Niel, G. W. Wardell-Johnson, C. J. Yates, M. Byrne, L. Mucina, A. G. T. Schut, S. D. Hopper, and S. E. Franklin. 2012. Refugia: identifying and understanding safe havens for biodiversity under climate change. *Global Ecology and Biogeography* 21:393–404.
- Kharouba, H. M., A. C. Algar, and J. T. Kerr. 2009. Historically calibrated predictions of butterfly species’ range shift using global change as a pseudo-experiment. *Ecology* 90:2213–2222.
- Kharouba, H. M., J. L. McCune, W. Thuiller, and B. Huntley. 2013. Do ecological differences between taxonomic groups influence the relationship between species’ distributions and climate? A global meta-analysis using species distribution models. *Ecography* 36:657–664.
- Kingston, D. G., M. C. Todd, R. G. Taylor, J. R. Thompson, and N. W. Arnell. 2009. Uncertainty in the estimation of potential evapotranspiration under climate change. *Geophysical Research Letters* 36:L20403.
- Klein, E., E. E. Berg, and R. Dial. 2005. Wetland drying and succession across the Kenai Peninsula Lowlands, south-central Alaska. *Canadian Journal of Forest Research* 35:1931–1941.
- Langsrud, Ø. 2003. ANOVA for unbalanced data: use Type II instead of Type III sums of squares. *Statistics and Computing* 13:163–167.
- Lawler, J. J., D. White, R. P. Neilson, and A. R. Blaustein. 2006. Predicting climate-induced range shifts: model differences and model reliability. *Global Change Biology* 12:1568–1584.
- Leroux, S. J., F. K. A. Schmiegelow, R. B. Lessard, and S. G. Cumming. 2007. Minimum dynamic reserves: A framework for determining reserve size in ecosystems structured by large disturbances. *Biological Conservation* 138:464–473.
- Lovette, I. J. 2005. Glacial cycles and the tempo of avian speciation. *Trends in Ecology and Evolution* 20:57–58.
- Lovette, I. J., and W. M. Hochachka. 2006. Simultaneous effects of phylogenetic niche conservatism and competition on avian community structure. *Ecology* 87(Supplement):S14–S28.
- Machtans, C. S., K. J. Kardynal, and P. A. Smith. 2014. How well do regional or national breeding bird survey data predict songbird trends at an intact boreal site? *Avian Conservation and Ecology* 9:5.
- Masson, D., and R. Knutti. 2011. Climate model genealogy. *Geophysical Research Letters* 38:L08703.
- Matthews, S. N., L. R. Iverson, A. M. Prasad, and M. P. Peters. 2011. Changes in potential habitat of 147 North American breeding bird species in response to redistribution of trees and climate following predicted climate change. *Ecography* 34:933–945.
- Mbogga, M. S., X. Wang, and A. Hamann. 2010. Bioclimate envelope model predictions for natural resource management: dealing with uncertainty. *Journal of Applied Ecology* 47:731–740.
- McKenney, D. W., M. F. Hutchinson, J. L. Kesteven, and L. A. Venier. 2001. Canada’s plant hardiness zones revisited using modern climate interpolation techniques. *Canadian Journal of Plant Science* 81:129–143.
- McKenney, D. W., J. H. Pedlar, K. Lawrence, K. Campbell, and M. F. Hutchinson. 2007. Beyond traditional hardiness zones: Using climate envelopes to map plant range limits. *BioScience* 57:929–937.
- McPherson, J. M., and W. Jetz. 2007. Effects of species’ ecology on the accuracy of distribution models. *Ecography* 30:135–151.
- McPherson, J. M., W. Jetz, and D. J. Rogers. 2004. The effects of species’ range sizes on the accuracy of distribution models: ecological phenomenon or statistical artefact? *Journal of Applied Ecology* 41:811–823.
- Meehl, G. A., C. Covey, T. Delworth, M. Latif, B. McAvaney, J. F. B. Mitchell, R. J. Stouffer, and K. E. Taylor. 2007. The

- WCRP CMIP3 multi-model dataset: a new era in climate change research. *Bulletin of the American Meteorological Society* 88:1383–1394.
- Mehta, V., G. Meehl, L. Goddard, J. Knight, A. Kumar, M. Latif, T. Lee, A. Rosati, and D. Stammer. 2010. Decadal climate predictability and prediction: Where are we? *Bulletin of the American Meteorological Society* 92:637–640.
- Mengel, R. M. 1964. The probable history of species formation in some northern wood warblers (Parulidae). *Living Bird* 3:9–43.
- Meynard, C. N., A. Migeon, and M. Navajas. 2013. Uncertainties in predicting species distributions under climate change: a case study using *Tetranychus evansi* (Acari: Tetranychidae), a widespread agricultural pest. *PLoS One* 8:e66445.
- Michaelian, M., E. H. Hogg, R. J. Hall, and E. Arsenault. 2010. Massive mortality of aspen following severe drought along the southern edge of the Canadian boreal forest. *Global Change Biology* 17:2084–2094.
- Murphy, J. M., D. M. H. Sexton, D. N. Barnett, G. S. Jones, M. J. Webb, M. Collins, and D. A. Stainforth. 2004. Quantification of modelling uncertainties in a large ensemble of climate change simulations. *Nature* 430:768–772.
- NABCI (North American Bird Conservation Initiative Canada). 2012. *The State of Canada's Birds, 2012*. Environment Canada, Ottawa, Canada.
- National Audubon Society. 2014. Audubon's birds and climate change report. <http://climate.audubon.org>
- Nielsen, S., D. Haughland, E. Bayne, and J. Schieck. 2009. Capacity of large-scale, long-term biodiversity monitoring programmes to detect trends in species prevalence. *Biodiversity and Conservation* 18:2961–2978.
- Noss, R. F. 2001. Beyond Kyoto: Forest management in a time of rapid climate change. *Conservation Biology* 15:578–590.
- Oliver, T. H., R. J. Smithers, S. Bailey, C. A. Walmsley, and K. Watts. 2012. A decision framework for considering climate change adaptation in biodiversity conservation planning. *Journal of Applied Ecology* 49:1247–1255.
- Pasher, J., E. Seed, and J. Duffe. 2013. Development of boreal ecosystem anthropogenic disturbance layers for Canada based on 2008 to 2010 Landsat imagery. *Canadian Journal of Remote Sensing* 39:42–58.
- Peng, C. H., Z. H. Ma, X. D. Lei, Q. Zhu, H. Chen, W. F. Wang, S. Liu, W. H. Li, X. Fang, and X. Zhou. 2011. A drought-induced pervasive increase in tree mortality across Canada's boreal forests. *Nature Climate Change* 1:467–471.
- Pennell, C., and T. Reichler. 2011. On the effective number of climate models. *Journal of Climate* 24:2358–2367.
- Peterson, A. T. 2001. Predicting species' geographic distributions based on ecological niche modeling. *Condor* 103:599–605.
- Peterson, A. T., M. A. Ortega-Huerta, J. Bartley, V. Sanchez-Cordero, J. Soberón, R. H. Buddemeier, and D. R. B. Stockwell. 2002. Future projections for Mexican faunas under global climate change scenarios. *Nature* 41:626–629.
- Podur, J., D. L. Martell, and K. Knight. 2002. Statistical quality control analysis of forest fire activity in Canada. *Canadian Journal of Forest Research* 32:195–205.
- Prentice, I. C., J. Bartlein, and W. Thompson. 1991. Vegetation and climate change in eastern North America since the last glacial maximum. *Ecology* 72:2038–2056.
- Price, D. T., et al. 2013. Anticipating the consequences of climate change for Canada's boreal forest ecosystems. *Environmental Reviews* 21:322–365.
- R Development Core Team. 2012. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Radeloff, V. C., et al. 2011. Economic-based projections of future land use in the conterminous United States under alternative policy scenarios. *Ecological Applications* 22:1036–1049.
- Räisänen, J. 2007. How reliable are climate models? *Tellus A* 59:2–29.
- Ralph, C. J., J. R. Sauer, and S. Droege. 1995. *Monitoring bird populations by point count*. USDA Forest Service General Technical Report PSW-GTR-149, Pacific Southwest Research Station, Albany, California, USA.
- Raupach, M. R., G. Marland, P. Ciais, C. Le Quéré, J. G. Canadell, G. Klepper, and C. B. Field. 2007. Global and regional drivers of accelerating CO₂ emissions. *Proceedings of the National Academy of Sciences USA* 104:10288–10293.
- Regan, H. M., Y. Ben-Haim, B. Langford, W. G. Wilson, P. Lundberg, S. J. Andelman, and M. A. Burgman. 2005. Robust decision making under severe uncertainty for conservation management. *Ecological Applications* 15:1471–1477.
- Rehfeldt, G. E., N. L. Crookston, C. Sáenz-Romero, and E. M. Campbell. 2012. North American vegetation model for land-use planning in a changing climate: a solution to large classification problems. *Ecological Applications* 22:119–141.
- Rich, T. D., et al. 2004. *Partners in Flight North American Landbird Conservation Plan*. Cornell Lab of Ornithology, Ithaca, New York, USA.
- Ridgely, R. S., T. F. Allnutt, T. Brooks, D. K. McNicol, D. W. Mehlman, B. E. Young, and J. R. Zook. 2005. Digital distribution maps of the birds of the western hemisphere. Version 2.1. NatureServe, Arlington, Virginia, USA.
- Ridgeway, G. 2012. Package gbm: generalized boosted regression models. Version 1.6.3.2. R Foundation for Statistical Computing, Vienna, Austria. <http://cran.r-project.org/web/packages/gbm/index.html>
- Roberts, D. R., and A. Hamann. 2012. Predicting potential climate change impacts with bioclimate envelope models: a palaeoecological perspective. *Global Ecology and Biogeography* 21:121–133.
- Rohling, E. J., et al. 2012. Making sense of palaeoclimate sensitivity. *Nature* 491:683–691.
- Rubidge, E. M., W. B. Monahan, J. L. Parra, S. E. Cameron, and J. S. Brashares. 2011. The role of climate, habitat, and species co-occurrence as drivers of change in small mammal distributions over the past century. *Global Change Biology* 17:696–708.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2011. *The North American Breeding Bird Survey, results and analysis 1966–2010*. Version 5.15.2011. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.
- Scherrer, S. C. 2011. Present-day interannual variability of surface climate in CMIP3 models and its relation to future warming. *International Journal of Climatology* 31:1518–1529.
- Schieck, J., and S. J. Song. 2006. Changes in bird communities throughout succession following fire and harvest in boreal forests of western North America: literature review and meta-analyses. *Canadian Journal of Forest Research* 36:1299–1318.
- Schmiegelow, F. K. A., C. S. Machtans, and S. J. Hannon. 1997. Are boreal birds resilient to forest fragmentation? An experimental study of short-term community responses. *Ecology* 78:1914–1932.
- Schneider, R. R., A. Hamann, D. Farr, X. Wang, and S. Boutin. 2009. Potential effects of climate change on ecosystem distribution in Alberta. *Canadian Journal of Forest Research* 39:1001–1010.
- Schneider, R. R., J. B. Stelfox, S. Boutin, and S. Wasel. 2003. Managing the cumulative impacts of land uses in the Western Canadian Sedimentary Basin: a modeling approach. *Conservation Ecology* 7:8.
- Screen, J. A., and I. Simmonds. 2010. The central role of diminishing sea ice in recent Arctic temperature amplification. *Nature* 464:1334–1337.

- Segurado, P., and M. B. Araújo. 2004. An evaluation of methods for modelling species distributions. *Journal of Biogeography* 31:1555–1568.
- Shaw, M. R., K. Klausmeyer, D. R. Cameron, J. MacKenzie, and P. Roehrdanz. 2012. Economic costs of achieving current conservation goals in the future as climate changes. *Conservation Biology* 26:385–396.
- Siriwardena, G. M., H. Q. P. Crick, S. R. Baillie, and J. D. Wilson. 2000. Agricultural land-use and the spatial distribution of granivorous lowland farmland birds. *Ecography* 23:702–719.
- Soberón, J. 2007. Grinnellian and Eltonian niches and geographic distributions of species. *Ecology Letters* 10:1115–1123.
- Soja, A. J., N. M. Tchebakova, N. H. F. French, M. D. Flannigan, H. H. Shugart, B. J. Stocks, A. I. Sukhinin, E. I. Parfenova, F. S. Chapin, III, and P. W. Stackhouse, Jr. 2007. Climate-induced boreal forest change: Predictions versus current observations. *Global and Planetary Change* 56:274–296.
- Sólymos, P., S. M. Matsuoka, E. M. Bayne, S. R. Lele, P. Fontaine, S. G. Cumming, D. Stralberg, F. K. A. Schmiegelow, and S. J. Song. 2013. Calibrating indices of avian density from non-standardized survey data: making the most of a messy situation. *Methods in Ecology and Evolution* 4:1047–1058.
- Stockwell, D. R. B., and A. T. Peterson. 2002. Effects of sample size on accuracy of species distribution models. *Ecological Modelling* 148:1–13.
- Stralberg, D., D. Jongsomjit, C. A. Howell, M. A. Snyder, J. D. Alexander, J. A. Wiens, and T. L. Root. 2009. Re-shuffling of species with climate disruption: a no-analog future for California birds? *PLoS One* 4:e6825.
- Strengers, B., R. Leemans, B. Eickhout, B. de Vries, and L. Bouwman. 2004. The land-use projections and resulting emissions in the IPCC SRES scenarios as simulated by the IMAGE 2.2 model. *GeoJournal* 61:381–393.
- Svenning, J.-C., and B. Sandel. 2013. Disequilibrium vegetation dynamics under future climate change. *American Journal of Botany* 100:1266–1286.
- Swanson, A. K., S. Z. Dobrowski, A. O. Finley, J. H. Thorne, and M. K. Schwartz. 2012. Spatial regression methods capture prediction uncertainty in species distribution model projections through time. *Global Ecology and Biogeography* 22:242–251.
- Synes, N. W., and P. E. Osborne. 2011. Choice of predictor variables as a source of uncertainty in continental-scale species distribution modelling under climate change. *Global Ecology and Biogeography* 20:904–914.
- Tebaldi, C., R. L. Smith, D. Nychka, and L. O. Mearns. 2005. Quantifying uncertainty in projections of regional climate change: a Bayesian approach to the analysis of multimodel ensembles. *Journal of Climate* 18:1524–1540.
- Thomas, C. D., and J. L. Lennon. 1999. Birds extend their ranges northwards. *Nature* 399:213.
- Thuiller, W. 2004. Patterns and uncertainties of species' range shifts under climate change. *Global Change Biology* 10:2020–2027.
- Thuiller, W., B. Lafourcade, R. Engler, and M. B. Araújo. 2009. BIOMOD—a platform for ensemble forecasting of species distributions. *Ecography* 32:369–373.
- Thuiller, W., S. Lavorel, M. B. Araújo, M. T. Sykes, and I. C. Prentice. 2005. Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences USA* 102:8245–8250.
- Veloz, S. D., N. Nur, L. Salas, D. Jongsomjit, J. Wood, D. Stralberg, and G. Ballard. 2013. Modeling climate change impacts on tidal marsh birds: Restoration and conservation planning in the face of uncertainty. *Ecosphere* 4:art49.
- Wang, M., J. E. Overland, V. Kattsov, J. E. Walsh, X. Zhang, and T. Pavlova. 2007. Intrinsic versus forced variation in coupled climate model simulations over the Arctic during the twentieth century. *Journal of Climate* 20:1093–1107.
- Wang, T., A. Hamann, D. L. Spittlehouse, and T. Q. Murdock. 2012. ClimateWNA—high-resolution spatial climate data for western North America. *Journal of Applied Meteorology and Climatology* 51:16–29.
- Weir, J. T., and D. Schluter. 2004. Ice sheets promote speciation in boreal birds. *Philosophical Transactions of the Royal Society B* 271:1881–1887.
- Wells, J., and P. Blancher. 2011. Global role for sustaining bird populations. Pages 7–22 in J. V. Wells, editor. *Boreal birds of North America: a hemispheric view of their conservation links and significance*. *Studies in Avian Biology* 41. University of California Press, Berkeley, California, USA.
- Wiens, J. J., et al. 2010. Niche conservatism as an emerging principle in ecology and conservation biology. *Ecology Letters* 13:1310–1324.
- Williams, J. W., and S. T. Jackson. 2007. Novel climates, no-analog communities, and ecological surprises. *Frontiers in Ecology and the Environment* 5:475–482.
- Williams, J. W., S. T. Jackson, and J. E. Kutzbach. 2007. Projected distributions of novel and disappearing climates by 2100 AD. *Proceedings of the National Academy of Sciences USA* 104:5738–5742.
- Wisz, M. S., R. J. Hijmans, J. Li, A. T. Peterson, C. H. Graham, and A. Guisan. 2008. Effects of sample size on the performance of species distribution models. *Diversity and Distributions* 14:763–773.
- Zurell, D., J. Elith, and B. Schröder. 2012. Predicting to new environments: tools for visualizing model behaviour and impacts on mapped distributions. *Diversity and Distributions* 18:628–634.
- Zurell, D., F. Jeltsch, C. F. Dormann, and B. Schröder. 2009. Static species distribution models in dynamically changing systems: how good can predictions really be? *Ecography* 32:733–744.

SUPPLEMENTAL MATERIAL

Ecological Archives

Appendices A–D and the Supplement are available online: <http://dx.doi.org/10.1890/13-2289.1.sm>